



# The Transgenerational Transmission of Trauma: The Effects of Maternal PTSD in Mother-Infant Interactions

Elisabetta Dozio, Marion Feldman, Cécile Bizouerne, Elise Drain, Mathilde Laroche Joubert, Malika Mansouri, Marie Rose Moro, Lisa Ouss

## ► To cite this version:

Elisabetta Dozio, Marion Feldman, Cécile Bizouerne, Elise Drain, Mathilde Laroche Joubert, et al.. The Transgenerational Transmission of Trauma: The Effects of Maternal PTSD in Mother-Infant Interactions. *Frontiers in Psychiatry*, 2020, 11, pp.480690. 10.3389/fpsy.2020.480690 . hal-03131921

**HAL Id: hal-03131921**

**<https://hal.parisnanterre.fr/hal-03131921>**

Submitted on 4 Feb 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# The Transgenerational Transmission of Trauma: The Effects of Maternal PTSD in Mother-Infant Interactions

Elisabetta Dozio<sup>1\*</sup>, Marion Feldman<sup>2</sup>, Cécile Bizouerne<sup>1</sup>, Elise Drain<sup>3</sup>,  
Mathilde Laroche Joubert<sup>2</sup>, Malika Mansouri<sup>4</sup>, Marie Rose Moro<sup>5</sup> and Lisa Ouss<sup>6</sup>

<sup>1</sup> Action Against Hunger, Paris, France, <sup>2</sup> Université Paris Nanterre, Nanterre, France, <sup>3</sup> Hôpital Avicenne, Bobigny, France, <sup>4</sup> Université Paris Descartes, Paris, France, <sup>5</sup> Hôpital Cochin, Paris, France, <sup>6</sup> Hôpital Necker-Enfants Malades, Assistance Publique Hôpitaux De Paris, Paris, France

## OPEN ACCESS

### Edited by:

Maria Muzik,  
University of Michigan, United States

### Reviewed by:

James Blair,  
National Institutes of Health (NIH),  
United States  
Jamie Lawler,  
Eastern Michigan University,  
United States  
Erika London Bocknek,  
Wayne State University, United States

### \*Correspondence:

Elisabetta Dozio  
bettadozio@hotmail.com

### Specialty section:

This article was submitted to  
Child and Adolescent Psychiatry,  
a section of the journal  
Frontiers in Psychiatry

Received: 25 June 2019

Accepted: 04 November 2020

Published: xx November 2020

### Citation:

Dozio E, Feldman M, Bizouerne C,  
Drain E, Laroche Joubert M,  
Mansouri M, Moro MR and Ouss L  
(2020) The Transgenerational  
Transmission of Trauma: The Effects of  
Maternal PTSD in Mother-Infant  
Interactions.  
Front. Psychiatry 11:480690.  
doi: 10.3389/fpsy.2020.480690

The objective of the study was to examine the process of mother to infant trauma transmission among traumatized mothers in humanitarian contexts. We investigated the impact of mothers' post-traumatic stress disorder symptoms on the quality of the dyadic interaction by conducting a microanalysis of mother-infant interactions at specific moments when trauma was recalled, compared to more neutral moments. Twenty-four mother-infant dyadic interactions of traumatized mothers and children aged from 1.5 to 30 months Central Africa, Chad, and Cameroon were videotaped during three sequences: a neutral initial session (baseline) exploring mothers' representations of the infant and of their bonding; a second sequence, "the traumatic narration," in which mothers were asked to talk about the difficult events they had experienced; and a third sequence focusing on a neutral subject. Three minutes of each sequence were coded through a specific grid for microanalysis [based on the scales developed at Bobigny Faculty of Medicine and the work of (1)], according to different communication modalities (touch, visual, and vocal), for both the mother and the child. Impact of traumatic event (IES-R), the level of depression and anxiety (HAD) were investigated in order to have a holistic understanding of the trauma transmission mechanism. The data analysis highlighted significant differences in mothers, children and their interaction during the "traumatic narration": mothers touched and looked at the infant less, looked more absent and smiled less, and looked less at the interviewer; infants looked less at the interviewer, and sucked the breast more. The mother-child interaction "infant self-stimulation—mother looks absent" and "Infant sucks the breast—mother looks absent" occurred more often during the mothers' traumatic narrations. The "absence" of the mother during trauma recall seems to have repercussions on infants' behavior and interaction; infants show coping strategies that are discussed. We found no significant associations between interaction and infant gender and age, the severity of traumatic experience, mothers' depression and anxiety symptoms, and the country of residence. The results of the microanalysis of interaction can shed light on the fundamental role of intermodal exchanges between mother and infant in trauma transmission during mothers' trauma reactivation.

**Keywords:** trauma transmission, mother-infant relationships, cross modal interactions, PTSD (post-traumatic stress disorder), infants

## INTRODUCTION

Trauma exposure is extremely common in countries affected by conflicts or natural disasters. In recent years, several studies have pointed to intergenerational trauma transmission (ITT) in countries affected by massive traumatic events (i.e., Burundi, Rwanda, Cambodia, Sierra Leone) and underlined the need to understand the mechanisms of transmission, in order to limit the potential negative impact on an entire community or even region, through generations (2–5).

Several studies in recent decades have tried to explain the transmission of parental trauma to the child. Many of these studies referred to collective historical trauma: the Holocaust, September 9/11, Vietnam Veterans, the Armenian genocide, etc., in which the impact of parents' post-traumatic stress disorder (PTSD) was measured in older children or in the second generation (6–10). Various other backgrounds have been explored in the process of ITT across generations, in different contexts of populations at risk for PTSD: low-income (11), abused mothers (12), adverse Childhood experiences (13) etc. Research has highlighted several possible models that may help explain the pathways of intergenerational trauma transmission.

Kellermann (14) proposed to distinguish the process of transmission, and what is transmitted. He described four models: psychodynamic, sociocultural, family system, and biological. Since this paper, two directions have been especially explored. Biological models have tried to understand the intergenerational transmission of stress (15) and the epigenetic mechanisms of stress and trauma transmission (16). Another set of studies concerns how the infants' sociocultural and family environment, the parents' representations, behavior and patterns of communication with their infant, can be considered as important factors involved in the process of ITT, a mechanism that appears to be a complex interaction between several co-processes [Perinatal Interactional Model of ITT, (17)]. Parental PTSD is characterized by physiological and emotional dysregulation, trauma related beliefs, and avoidance/withdrawal. This impacts fetal programming, and has repercussions on parent-child interaction through social learning, and maladaptive parenting. These phenomena decrease the child's regulatory capacity, increase child distress, and can lead to poorer child mental health, which in turn participates in a vicious circle.

The attachment theory (18) offers a theoretical framework to explain how caregiving behavior is impacted by parents' traumatic experience and can directly affect the child. Parents with unresolved trauma experience dissociation phenomena. They show frightened emotional expressions to the child who is unable to make the causal link between the terror of his/her parents and the trauma. The child responds with emotional and behavioral disorganization (19) and disorganized attachment behavior (20).

Maternal cognitions and representations also impact caregiving behavior, through reflective functioning. Schechter et al. (21) found that greater maternal PTSD severity was associated with unrealistic expectations of the child and distorted attributions of child intentions. These distorted maternal mental representations of the child (22, 23) will increase the frightening

behaviors in response to the child, who consequently will react with disorganized behaviors. Child distress and reactions can reactivate traumatic memories in the mother, leading to a vicious circle. This can lead to specific offspring attachment orientations, in particular to higher attachment insecurities, either anxiety or avoidance, in association to self-amplifying cycle of PTSD and attachment insecurities of parents following trauma exposure (24).

Emotional availability (25) focuses more specifically on the emotional signaling between mother and child as a major mediator of trauma transmission. High scores indicate adequate maternal emotion regulation, and the open expression of negative emotions results in lower scoring. It has been proposed as a process involved in ITT, in spite of the fact that it was not found as a mediator between maternal trauma and infant psychosocial functioning in a sample of refugee mothers (26).

The PTSD mechanism was confirmed in a recent longitudinal study of ITT, which suggested a strong effect of maternal PTSD on the attachment relationship and consequently on child development; however, the need to identify the mechanisms by which maternal PTSD has an impact on the mother-infant relationship was underlined (27). The interpersonal neurobiological model of attachment and relational trauma developed by Schore (28, 29) can help to explain maternal PTSD transmission to the infant through a mechanism involving the rhythmic pattern behind mother-infant interactions (visual, tactile, vocal), which are negatively affected when mothers are traumatized. Mothers cannot modulate their stimulation and their response to their infant's needs, leading to overstimulation or on the contrary, to neglect of the child. Both behaviors can be traumatic for the infant who will have difficulty self-regulating his affective state to protect himself from becoming overwhelmed by hyperstimulation or the absence of stimulation from the mother. In a refugee sample, mothers' intrusion and avoidance explained individual differences in extremely insensitive parenting, which had direct negative effects on children's attachment organization (26).

In line with this mechanism, through the concept of "Affective Attunement," Stern (30) proposed a pathway of transmission based on the sharing of emotional states between mother and child by means of inter-modal exchanges. D. Stern uses the term "intermodal" when in the dyadic exchanges an affect is expressed according to one modality (vocal, visual, bodily) then is "translated" and associated with another modality.

Affective attunement is a process centered on the mother's intermodal transformation of the baby's affective state. The mother tends to complement the baby's actions with gestures, gaze, and vocalizations, while the infant matches the mother's gesture behavior and perceives this as an affirmation of the continuity of his own affective state. The mother's transformation of the infant's affective state through behavior emphasizes the infant's recognition of his internal state. Traumatized mothers can transmit their negative affective states through their behavior, and these behaviors can modify the process of infant internalization of mothers' intentions.

More recently, a paradigm shift has been proposed for ITT in complex settings (war, migration). Intermediary dimensions

have been introduced between parental PTSD and their effects on parent-child relationships. When parents were less able to draw on secure attachment representations, symptoms of PTSD increased the risk of insensitive parenting in a sample of Dutch asylum seekers and refugee parents (31). It is difficult to isolate the contribution of mothers' Post-Traumatic Stress Disorder (PTSD) in ITT, as in most cases, similarities with depressed or anxious mothers are frequent.

Studies of the direct mechanisms behind the transmission of parental traumatic experience to young children through caregiving have been undertaken recently, but are still rare [(Schuengel, 1999; Schechter, 2008), (6, 26, 27, 32–35)].

To investigate trauma transmission in a population at high risk of trauma exposure, we chose population from Central African Republic (CAR), which has been affected by collective violence since 2013, forcing the population to live in internal displaced camps or to migrate to refugee camps in border countries such as Chad or Cameroon. The current study addresses the issue of psychological trauma of the mother, which occurred in her life before the child was born. The objective was to identify the direct specific processes of mother-to-infant trauma transmission in traumatized mothers in humanitarian contexts, through the comparison of inter-modal mother-child interactions, occurring during different paradigmatic moments: moments of trauma recall (possibly leading to dissociation), compared to neutral moments, using a microanalysis method. We hypothesized that mothers affected by maternal post-traumatic stress symptoms would show lower levels of availability to respond to their infants' needs and show poor or inappropriate mother-infant interactions. When faced with the traumatic experiences or their recall (36), mothers may show specific behaviors (absent gaze, disconnection from reality, etc. as manifestations of dissociative states), which will result in a lack of attunement in interactions: non-synchrony (temporal coordination) or non-qualitative contingency (lack of inter-modality) between mother and infant behaviors. We hypothesized that during the narration of their traumatic experience in presence of the child, mothers may experience episodes of dissociation, that impact the infant's reactions and the interactions between mother and infant. Observing this specific moment in comparison with a neutral moment could highlight the specific mechanisms involved in trauma transmission.

## METHODS

### Participants

Inclusion criteria: participants in the study were mother and child dyads in which the mothers had been exposed to traumatic events, according to criteria A of DSM V, in absence of the infant, or before the child's birth, including during pregnancy. Children were aged 1–36 months.

The recruitment took place in three African countries affected by the political crises in the Central African Republic (CAR), which started in 2013. In CAR, the sample comprised internally displaced mothers and their infants, whereas in Chad and Cameroon, the sample comprised refugees who had fled the ongoing violence in CAR.

The dyads were selected in accordance with the inclusion criteria by psychosocial workers working in the international NGO *Action Contre la Faim* (ACF) psychosocial support programs in the three countries. All the participants provided their written informed consent. Mothers were given the opportunity to ask questions to better understand the aims of the study before agreeing to take part and could refuse to continue the interview at any time.

The research protocol was approved by the institutional ethical review boards in CAR and in Cameroon, and by the "Ministère de la Femme, de l'Action Sociale et de la Solidarité Nationale" (Ministry of Women, Social Action and National Solidarity) in Chad.

### Procedure

The mothers were invited to participate in semi-structured interviews in the presence of their infant. The interview was videotaped to allow the microanalysis of mother and child behaviors and their interactions.

A clinical psychologist interviewed each mother with her infant in presence of the psychosocial worker who was in charge of the mother's psychological follow-up in the ACF program. This option was chosen in order to establish greater trust between the mother and the investigator and to guarantee a follow up in the case of increased distress in mothers and or infants during or after the research interview. The psychosocial worker was also the interpreter, as the investigator did not speak the local languages.

To investigate the core process of trauma transmission we developed a paradigm involving different settings, which explores the mother-infant interactions during three different moments: one moment of trauma recall, and two neutral moments, in order to compare within each dyad, each dyad being its own control, the changes in mother-child interaction.

- First sequence: a neutral initial session considered as the baseline to compare the other two sessions, "before the traumatic narration." Some neutral questions explored mothers' representations of the infant and of their bonding,
- Second sequence: "the traumatic narration." Mothers were asked to answer the question: "Could you tell me something about the difficult events you have experienced?" They were invited to narrate their traumatic events with the maximum of details (nature, temporality, numbers of events, etc.) and the effect on their present life (migration, change of situation, loss of family links, mourning, etc.). This was not the first time they narrated the events as they had already done so during the psychological follow up with the psychosocial worker.
- Third sequence: "after the traumatic narration," began with the first mention by the mother of a neutral subject (i.e., the naming of the child, her projection into the future, cultural practices surrounding birth, etc.). The aim of these questions was to stabilize the mothers' emotional state.

### Measures

*Impact of traumatic events and Post-traumatic Stress Disorder (PTSD) symptoms* of mothers were screened using the Impact of



Event Scale revised (IES-R) questionnaire (37). The scale consists of a list of 22 self-reported items assessing the perceived distress caused by traumatic events. Participants are asked to identify a specific stressful life event and then to say to what extent they had been affected during the past week on a 5-point Likert scale ranging from 0 “not at all” to 4 “extremely.” The IES-R total score ranges from 0 to 88 and subscale scores can also be calculated for the intrusion (8 items), avoidance (8 items), and hyperarousal (6 items) subscales. For each of these subscale scores, it is recommended to use means instead of raw sums. The maximum mean score on each of the three subscales is “4.” The IES-R is widely used for a preliminary diagnosis of PTSD (38). For the general IES-R score (calculated using summing), 33 and above represents the best cut-off for a probable diagnosis of PTSD (39). The literature shows that even though not formally validated in all the different contexts, the IES-R scale is used to measure PTSD symptoms in many cultures throughout the world (40). For this study, the French version of the IES-R was used, which has satisfactory internal validity and test-retest reliability (41).

*Symptoms of depression and anxiety* were measured using the Hospital Anxiety and Depression scale (HAD) (42). The HAD is an instrument for symptoms related to anxiety (HADS-A) and depression (HADS-D) composed of 14 items. Each question asks about the frequency of specific symptoms in the past week using a 4-point scale ranging from 0 (*not at all*) to 3 (*very often*). For each subscale, the cut-off identified is a point of 8/21. This instrument has been validated in both medical and general populations, and the results of a recent systematic review on screening tools for common mental disorders in low and middle income countries (43) broadly recommend using the HAD scale for depressive and anxiety disorders, as it been validated in multiple settings.

Microanalysis of Mother and Child Behaviors

We chose to conduct a microanalysis of the mother-child interaction in order to identify instant-by-instant events and the intermodality of interaction. Videotaped mother and child interactions during the interview were analyzed with an observation grid based on the Action Research Training (RAF, *Recherche Action Formation*) scales developed at the Bobigny Faculty of Medicine (44) and on the microanalysis conducted by Beebe et al. (45) on mother-infant interactions (Table 1). These previous studies analyzed mother/infant or mother/toddler behaviors according to the different types of communication: tactile, vocal and visual.

In addition to the distinct behavior and communication actions of mother and child, a list of possible communication overlaps, which could suggest a possible interaction, was defined. Based on the most frequently observed interactions, 28 mother-child communication modality pairs were retained for the analysis (Table 2).

We selected the first 3 min of each of the three moments of videotaped interview (before, during and after the “traumatic narration”) for analysis.

In each sequence of 3 min, any change in behavior, among those listed in the grid shown in Table 1, was noted and measured

TABLE 1 | Observation grid for mother and infant communication modalities.

Modality of communication	Behavior observed	
	Mother	Infant
Bodily	Touches the infant	Touches the mother
	Touches an object	Touches an object
Visual		Self-stimulation/ self-regulation
		Sucks the breast
		Sleeps
	Looks at the infant	Looks at the mother
	Look at what is doing the infant	Looks at the interviewer
	Looks at the interviewer	Look at the environment
Vocal	Look at the environment	Looks at an object
	Looks absent	Looks at his body
	Speaks to the infant	Looks absent
	Vocalizes to the infant	Vocalizes
Affects	Speaks to the interviewer	Vocalizes to the mother
	Smiles	Vocalizes to the interviewer
	Smiles at the infant	Smiles
	Cries	Smiles at the mother
		Cries

in terms of duration. Interrupts of the same action lasting <0.50 s were not taken into account. The action was then coded as continuous. “Touch” was coded as such only if the gesture was intentional. Contacts between mother and baby caused by unintentional gestures were not considered. The fact that the child was sitting on the mother’s lap was not taken into account in intentional tactile interactions.

The frequency and the duration of the mother-infant inter-modal interactions were coded using the open source multimedia annotator Elan, version 4.9.3 (46). The software was used to code the length of intermodal distinct behavior and interactions. Communication overlaps (see Table 2) were generated automatically by the Elan software. Inter-rater reliability (Cohen’s Kappa) of the coding of mother and infant behaviors was calculated on a randomly selected sample of 25% of the dyads interviewed and was satisfactory ( $k = 0.686$ ).

Statistical Analysis

Friedman’s test was used for the analysis of difference in behaviors and interactions before, during, and after the traumatic narration by the mother.

The choice of a non-parametric test was made because of the small sample size. Among the non-parametric tests, Friedman’s test is an ideal statistic to use for a repeated measures type of experiment to determine if a particular factor has an effect (47). The Friedman test is a non-parametric alternative to the repeated measures ANOVA when the assumption of normality is not acceptable.

In addition, to examine where the differences actually occurred, we ran separate Wilcoxon signed-rank tests on the different combinations of related groups. Spearman’s correlation was used to analyse correlations between scale scores and

**Q21** **TABLE 2 |** Grid for microanalysis of mother-infant interactions.

**Mother–infant interactions**

Infant touches the mother - Mother touches the infant
Infant touches the mother - Mother looks at the infant
Infant self-stimulation - Mother looks absent
Infant sucks the breast - Mother touches the infant
Infant sucks the breast - Mother looks at the infant
Infant sucks the breast - Mother looks absent
Infant looks at the mother - Mother touches the infant
Infant looks at the mother - Mother looks at the infant
Infant looks at the mother - Mother looks absent
Infant look at an object - Mother looks at what is doing the infant
Infant looks absent - Mother touches the infant
Infant looks absent - Mother looks at the infant
Infant looks absent - Mother looks absent
Infant vocalizes - Mother vocalizes to the infant
Infant vocalizes - Mother speaks to the infant
Infant vocalizes - Mother touches the infant
Infant vocalizes - Mother looks at the infant
Infant vocalizes - Mother looks absent
Infant smiles - Mother touches the infant
Infant smiles - Mother looks at the infant
Infant smiles - Mother looks absent
Infant smiles at the mother - Mother miles at the infant
Infant cries - Mother touches the infant
Infant cries - Mother looks at the infant
Infant cries - Mother looks absent
Infant cries - Mother smiles

demographic variables. All statistical analyses were performed using SPSS version 21.

**RESULTS**

**Descriptive Results**

Twenty-four mother and infant dyads met the inclusion criteria of this study. Sixteen were recruited in CAR, three in Chad and five in Cameroon. The demographic data for the 24 dyads are presented in **Table 3**.

The children’s age ranged from 1.2 to 30 months [ $M = 11.94$ ,  $SD = 6.92$ ; Median = 10.55, Interquartile Range (IRQ) = 6.3] and consisted of 11 boys and 13 girls.

The mothers were aged between 16 and 37 [ $M = 26$ ,  $SD = 16.18$ ; Median = 27, Interquartile Range (IRQ) = 11]. Some mothers were unable to indicate their own age, as they did not know it. This is quite frequent in the rural contexts of countries in the field of this study. Eighty-three percent of the mothers

were pregnant during the traumatic event; four children were born during traumatic situations. Only one mother was pregnant during the interview. At the time of the interview, all of the children were the youngest of the siblings. The father was present in 62.5% of the cases. His absence was due either to his death or to his disappearance, without having given any news.

Mothers had experienced multiple traumatic events. The detail for each mother is shown in **Table 3**: among the traumatic events they had witnessed the murder of their children, or husband, relatives or friends, some of them witnessed massive violence in the community and/or had been forced to leave their homes, lost all their possessions, and feared for themselves and their loved ones, etc. The time between the traumatic event and the interview was on average 16 months, with a minimum of 7 and a maximum of 31 months (Median = 13, IRQ = 11).

As shown in **Table 4**, the impact of the traumatic event was very high among mothers. In addition, they presented high rates of depression and anxiety.

**Correlations Between Clinical Status and Demographic Variables**

Correlations were run to assess the relationships between the clinical status (scores of HAD and IES-R scales in **Table 4**) and the main demographic variables reported in **Table 3**. Spearman’s correlation coefficient was used rather than the Pearson correlation coefficient because of the small sample size ( $n = 24$ ).

No correlation was found between scores at HAD or IES-R and the age of the mother, the gender and age of the child, the presence of the father, the fact of being primiparous and the number of children.

Results of the Spearman correlation indicated that there was a significant positive association between the severity of the impact of the traumatic event (IES-R) and the time that had elapsed since the traumatic events ( $r = 0.54$ ,  $p = 0.01$ ), meaning that the more time had passed since exposure to the traumatic event, the more the participants showed traumatic symptoms, in particular for intrusion ( $r = 0.45$ ,  $p = 0.05$ ) and hyper arousal ( $r = 0.468$ ,  $p < 0.001$ ), both measured through IES-R subscales.

The Wilcoxon-Mann-Whitney test was used to cross the qualitative variables with two classes such as the gender of the child, with the quantitative variable of the clinical scales. No significant differences were found between the gender of the child and the clinical profile of the mother.

A Kruskal-Wallis test was conducted to examine the differences in mothers’ clinical symptoms according to the country of residence. Results indicated that the highest level of PTSD symptoms was associated with populations still living in CAR,  $H(2) = 8.5$ ,  $p = 0.014$ .

No other clinical symptoms (Anxiety and Depression) were linked to the Country of residence.

**Results of Microanalysis**

Results refer to separate actions by the mother and by the infant, and to their interactions.

**TABLE 3 |** Demographic data of participants.

Dyad	Country	Age of the mother (years)	Age of the child (months)	Sex of the child	Number of children in the family	Type of traumatic event	Mother : pregnant during the traumatic event	Presence of the father	Raison of the absence of father
1	Cameroon	35	11	F	7	1;2;4;5	Yes	Yes	–
2	Cameroon	30	7	F	3	1;2;3;4;8	Yes	Yes	–
3	Cameroon	25	1,5	F	6	1;2;3;4	No	Yes	–
4	Cameroon	26	12	F	MD*	1;2;3;5	Yes	Yes	–
5	Cameroon	MD	6	F	MD	1;2;3;6;7	Yes	No	Disappeared
6	CAR	18	9	M	1	1;3;5;7	Yes	No	Disappeared
7	CAR	28	30	M	5	1;3;4	Yes	Yes	–
8	CAR	37	7	F	4	1;3;4;8	Yes	Yes	–
9	CAR	MD	17	M	1	1;3;4;6	No	Yes	–
10	CAR	MD	17	M	?	1;3;4	Yes	Yes	–
11	CAR	19	8	F	1	1;3;4;7	Yes	No	Disappeared
12	CAR	MD	12	F	MD	1;3;4;6	Yes	No	Dead
13	CAR	16	12	M	1	1;3;4	Yes	Yes	–
14	CAR	26	11	M	2	1;3;6	Yes	Yes	–
15	CAR	31	24	M	3	1;3;4	Yes	Yes	–
16	CAR	28	14	F	3	1;3;4;5;6	Yes	Yes	–
17	CAR	16	7	F	1	1;3	Yes	Yes	–
18	CAR	20	8	F	2	1;3;4;5;6;7	No	No	Disappeared
19	CAR	28	7	M	2	1;3;4	Yes	Yes	–
20	CAR	29	28	M	9	1;3;4	Yes	Yes	–
21	CAR	32	10	M	6	1;3;4;5	Yes	No	Dead
22	Chad	MD	12	F	3	1;3;4	No	No	He stayed in CAR
23	Chad	MD	8	F	MD	1;3;4	Yes	No	He Fled
24	Chad	25	8	M	1	1;3;4;5	Yes	No	Dead

\*MD, Missing Data.

1 Escape from gunfire, detonations, violence.

2 Migration on foot in extremely difficult conditions.

3 Exposure to combat and violence.

4 Witness to the murders.

5 Witness to the death of a family member.

6 Death of loved one due to the conflict.

7 Husband's disappearance.

8 Death of his own child(ren).

**TABLE 4 |** Clinical status.

Measure	Outcome	Min	Max	M	SD	Median	IRQ	Range
HAD	Anxiety	5	18	13.08	3.425	14	4.5	Clinical
	Depression	8	17	12.33	2.180	12	3	Clinical
IES-R	Avoidance	1	3	1.96	0.374	1.85	0.7	High
	Intrusion	1	3	2.36	0.687	2.55	2.6	High
	Hyperarousal	0	3	2.22	0.817	2.25	1.2	High
	Total PTSD	20	64	47.08	10.685	49.5	12.8	Clinical

### Differences in the Mothers' Behaviors Between the Three Sessions

Analysis revealed the following statistically significant differences: the mother spoke to the interviewer ( $p = 0.030$ ) more during the traumatic narration than in the session after ( $Z = -2.171$ ,  $p = 0.030$ ); the mother touched the infant ( $p = 0.001$ )

less during the traumatic narration than in the session before ( $Z = -3.589$ ,  $p = 0.000$ ) and more than in the session after ( $Z = -2.886$ ,  $p = 0.004$ ); the mother looked at the infant ( $p = 0.018$ ) less during the traumatic narration than after ( $Z = -3.494$ ,  $p = 0.000$ ); the mother looked at the interviewer ( $p = 0.006$ ) less during the traumatic narration than before

685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741

**Q21** **TABLE 5 |** Differences in the duration of the mothers' behaviors in the three sessions.

Q22	3 min session				3 min session				3 min session				Asymp	Pairwise comparison	
Q23	before (a)				during (b)				after (c)						
	Min	Max	Mean (SD)	MD (interquartile range IQR)	Min	Max	Mean (SD)	MD (IQR)	Min	Max	Mean (SD)	MD (IQR)			(Wilcoxon sing-ranked test)
Vocal interactions															
Mother speaks to the infant	0	1.1	0.08 (0.28)	0 (0)	0.8	2.3	0.18 (0.55)	0 (0)	0.3	16.6	1.37 (3.46)	0 (1.9)	0.139		
Mother vocalizes to the infant	0	6.3	2.61 (1.27)	0 (0)	0	0	0 (0)	0 (0)	0	0	0 (0)	0 (0)	0.368		
Mother speaks to the interviewer	26.4	136.7	65.23 (28.43)	58.2 (35.7)	30.03	180.7	75.02 (35.25)	63.8 (22.9)	0	104.6	55.85 (25.79)	53.1 (36.6)	0.030*	b > a; b > c*	
Touch interactions															
Mother touches the infant	0	152.7	58.34 (47.07)	46.1 (79.2)	0.1	118.3	23.35 (28.71)	14.7 (-23)	0.2	164.2	52.13 (40.46)	47.2 (67.6)	0.001*	b < a*; b < c*	
Mother looks at the infant	0	20.03	6.57 (6.83)	3.9 (13.4)	0	14.7	3.35 (4.45)	1.6 (5.7)	0	45.1	10.68 (11.79)	7.1 (9.5)	0.018*	b < a; b < c*	
Mother looks at what the infant is doing	0	46.9	8.69 (10.93)	4.9 (13.2)	0	20.6	3.66 (5.31)	1.7 (5.9)	0	39.8	6.75 (10.26)	2.2 (9)	0.096		
Mother looks at the interviewer	23.8	178.5	114.95 (28.12)	11.5 (48.3)	19.7	169.6	90.69 (39.53)	91 (57.1)	22.9	156.9	106.82 (37.43)	115.6 (55.6)	0.006*	b < a*; b < c*	
Mother looks at the environment	0	73.1	14.06 (15.55)	9.9 (13.3)	0	98.3	18.37 (26.4)	8.2 (19.5)	0	107.9	18.07 (23.61)	9.8 (16)	0.153		
Mother looks absent	0	145	34.78 (37.43)	22.3 (34.1)	0	146	59.17 (44.1)	50.9 (74.6)	0	119	36.8 (32.12)	25.6 (35.3)	0.004*	b > a*; b > c*	
Affects															
Mother smiles	0	57.9	13.0 (18.26)	4.04 (17.4)	0	47.6	3.59 (10.63)	0 (0.5)	0	77	12.15 (21.13)	3.25 (16.6)	0.004*	b < a*; b < c*	
Mother cries	0	0	0 (0)	0 (0)	0	88.2	3.67 (18)	0 (0)	0	0	0 (0)	0 (0)	0.368		

\*p < 0.05.

**Q21** **TABLE 6 |** Differences in the duration of infants' behaviors in the three sessions.

Q22

Q24

	3 min session before (a)				3 min session during (b)				3 min session after (c)				Asymp Sig.	Pairwise comparison (Wilcoxon sing-ranked test)
	Min	Max	Mean (SD)	MD (interquartile range IQR)	Min	Max	Mean (SD)	MD (IQR)	Min	Max	Mean (SD)	MD (IQR)		
Vocal interaction														
Infant vocalizes	0	69.2	6.24 (14.94)	1.51 (5.1)	0	70.3	10.85 (18.35)	0.85 (11.4)	0	35.1	7.29 (10.09)	1.25 (13.3)	0.532	
Touch interactions														
Infant touches the mother	0	98.7	27 (30.26)	11.51 (45.2)	0	174.3	44.21 (43.87)	31.15 (55.1)	0	167.4	46.2 (49.03)	28.8 (49.3)	0.158	
Infant touches an object	0	180.4	95.73 (68.88)	88.49 (145.5)	0	180.7	83.61 (71.66)	82.5 (152.6)	0	180.8	58.21 (66.89)	31.5 (123.4)	0.071	
Infant self-stimulation	0	179.7	21.18 (40.29)	4.03 (28.8)	0	151.2	29.45 (41.31)	9 (38.9)	0	173.1	25.21 (498)	2.65 (27.5)	0.542	
Infant sucks the breast	0	180.0	16.45 (42.76)	0 (0)	0	180.8	50.72 (71.58)	0 (100.3)	0	152.8	27.81 (51.53)	0 (42)	0.045*	b > a*; b > c
Visual interactions														
Infant looks at the mother	0	29.1	2.47 (6.12)	0.2 (1.7)	0	10.2	1.28 (2.5)	0 (1.6)	0	15.2	3.7 5 (4.71)	1.65 (5.7)	0.288	
Infant looks at the environment	0	131	39.15 (39.37)	31.8 (56)	0	122.2	43.42 (34.39)	42.85 (44.9)	0	104.2	36.32 (30.72)	27.9 (62.8)	0.722	
Infant looks at the interviewer	0	103.7	37.34 (27.94)	30.29 (31.2)	0	79.2	16.45 (20.56)	7.3 (24.3)	0	71.9	16.23 (20.1)	10.2 (26.5)	0.001*	b < a*; b > c
Infant looks at his own body	0	28.5	3.25 (6.93)	0 (3.3)	0	20	1.76 (4.39)	0 (1.9)	0	41.8	2.5 (8.7)	0 (0)	0.195	
Infant looks absent	0	64.4	24.78 (21.04)	20.4 (38.1)	0	86.9	24.78 (30.16)	10.25 (43)	0	78.9	18.59 (26.46)	3.15 (30.8)	0.364	
Affects														
Infant smiles	0	7.5	1.28 (2.25)	0 (3.2)	0	3.5	0.73 (1.17)	0 (1.4)	0	19.9	1.85 (20.06)	0 (2)	0.980	
Infant cries	0	18.1	1.8 (4.97)	0 (0)	0	57.3	4.99 (12.92)	0 (1.2)	0	86.5	8.4 (20.5)	0 (1)	0.412	

\*p < 0.05.

742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798



( $Z = -2.886$ ,  $p = 0.004$ ) and less than in the session after ( $Z = -2.800$ ,  $p = 0.005$ ); the mother looked absent ( $p = 0.004$ ) more during the traumatic narration than before ( $Z = -2.950$ ,  $p = 0.003$ ) and more than after ( $Z = -3.346$ ,  $p = 0.001$ ); the mother smiled ( $p = 0.004$ ) less during the traumatic narration than before ( $Z = -2.675$ ,  $p = 0.007$ ) and after ( $Z = -2.045$ ,  $p = 0.041$ ). No other significant differences were found.

### Qualitative Analysis of Mothers' Behavior

The mothers' vocal interactions with the infant were very limited; mothers were more involved in speaking with the interviewer, especially when they were recounting the traumatic event. The mothers' visual interactions with the infant were poor and more focused on the interviewer.

All the mothers touched the infant at least once during one of the three sessions analyzed. Most of the time during the interview the mother showed a neutral facial affect except for a visible change of gaze when narrating the traumatic event. Only one mother cried and 17 mothers smiled at least once, but they never smiled at the infant.

### Differences in the Infants' Behaviors Between the Three Sessions

The repeated measures compared using Friedman's test revealed a statistically significant difference in: "Infant looks at the interviewer" ( $p = 0.001$ ) and "Infant sucks the breast" ( $p = 0.045$ ). *Post-hoc* analysis with Wilcoxon signed-rank tests was conducted and showed that the infant looked at the interviewer less during the mother's traumatic narration than in the session before ( $Z = -3.360$ ,  $p = 0.001$ ) and they sucked the breast more during the mother's traumatic narration than before ( $Z = -2.395$ ,  $p = 0.017$ ). No other changes in infant behavior were significant in the three sessions.

### Qualitative Analysis of Infants' Behavior

As with the mothers, very few *vocal* infant interactions were observed during the sessions. Vocalizations were generally neutral or positive. To determine the direction of the infant's vocalization, we observed the direction of gaze and / or the spatial position of the infant's body, to understand if the vocal interaction was toward the mother and/or the interviewer. Infants produced no vocalizations toward the mother or the interviewer.

The infants' *visual* interactions tended to be directed toward an object: a toy, the chair where they were seated with their mother, objects on the ground, etc. When the infant changed the direction of his gaze, it was toward the interviewer or the environment, rather than toward the mother. Touch was the most frequent communicative means used by infants. All the infants touched their mother at least once during the three sessions. In particular, the tactile interaction was more frequent during the traumatic narrative. The majority of children (21 of 24) showed self-stimulation behaviors. We included in self-stimulation broader behaviors represented by the repetition of physical movements (e.g., repetitive swaying, movement of the fingers, hands, etc.), which can be self-regulation behaviors.

Sucking the breast was included among the physical interactions and 11 infants suckled at least once during the three sessions. These infants were aged from 1.5 to 17 months ( $M = 11.5$ ,  $SD = 4.47$ ; Median = 12, IRQ = 4). In general, infants showed a neutral expression during the sessions. One third expressed affect by smiling or crying.

### Differences in the Mother-Infant Interactions Between the Three Sessions

The repeated measures of all the 28 interactions between mother and child revealed that only two variables were statistically significant: "Infant self-stimulation—Mother looks absent" ( $p = 0.045$ ) that occurred more during the mother's traumatic narration than in the session before ( $Z = -2.999$ ,  $p = 0.003$ ) and more than in the session after ( $Z = -2.731$ ,  $p = 0.006$ ). "Infant sucks the breast—Mother looks absent" ( $p = 0.003$ ) occurred more during the traumatic narration than before ( $Z = -2.090$ ,  $p = 0.037$ ) and more than after ( $Z = -2.118$ ,  $p = 0.034$ ).

Correlations were run to assess the associations between the mother-infant interactions and demographic variables such as age and gender of infant, age of the mother, presence of the father, etc., and clinical status of the mothers (HAD and IES-R scores). No significant associations were found.

### Qualitative Analysis of Mother-Infant Interactions

The analysis of mother and infant interactions showed very little reciprocity and contingency of interactions. Interaction in the same modality: "infant touch—mother touch," "infant vocalization—mother vocalization or speaking," "infant looks at the mother—mother looks at the infant," were rare.

## DISCUSSION

The purpose of this study was to shed light on the process of mother to child trauma transmission in mothers who had experienced severe traumas in humanitarian contexts. As far as we know, this is the only study to investigate the quality of the dyadic interaction, by observing and conducting a microanalysis of the mother-infant interactions in a trauma reactivation situation. The mother-child interaction during a trauma recall situation was compared with mother-child interaction during a more neutral interview.

We found three main results, confirming our hypothesis.

The first result was that the mothers' behavior toward the child was specifically affected during trauma recall narration compared to the two more neutral moments of interaction. During the "traumatic narration," mothers appeared to be emotionally affected by memories: they touched the infant less than in the session before, they looked at the infant less than in the session after, they looked more absent and smiled less than "before" and "after" moments. Moreover, they looked at the interviewer less during the traumatic narration than before and after. This is in favor of the relevance of our setting, which tried to catch these dissociative moments. When they were faced with the trauma recall, the mothers' absent look suggested that they were re-experiencing the event and were unable to focus on the present situation and on the infant's requests. They showed

a behavioral pattern linked to the traumatic experience: they withdrew from interaction, smiled, and looked less at others, and seemed to be absent. These signs may reflect the maternal traumatic dissociation the infants face when their mother is lost in the traumatic experience. The mothers seem cut off from the present experience without being able to perceive external reality and the needs expressed by their child (48). This suggests that the mothers had difficulty in properly assessing the verbal and non-verbal expressions of infant arousal because they were overwhelmed by their own emotional state. This “absence of an appropriate response” emphasizes the transmission of the mothers’ negative emotional state to the infants, who can potentially internalize this affective state as their own (30).

The majority of children in this study had no access to the direct meaning of their mother’s words as they still were in a pre-verbal developmental stage, and even for those aged between 12 and 30 months, their mothers’ words could not attack their symbolic world since they were not yet able to organize Internal Working Models, due to their developmental stage (49). Consequently, what is traumatic for them is more related to the absence, the lack of response to their requests and queries, which can potentially create external and internal sensory chaos (50).

The second finding is that the infants’ behavior was impacted by their mothers’ traumatic experience. During the traumatic narration, infants looked at the interviewer less than in the session before, and they sucked the breast more than before. Moreover, the mother-child interaction was also affected: “Infant self-stimulation—Mother looks absent” and “Infant sucks the breast—Mother looks absent” occurred more during the traumatic narration than in the other two sessions. These interactive behaviors reflect the children’s coping strategies of self-regulation: they perceive these moments, and try to compensate with self-stimulation or by clinging to the satisfaction of primary needs. To suck the breast is a behavior which maintains close contact with the mother, and offers a primary satisfaction that the child does not find in the interaction. The lack of correlation between the age of the child and suckling the breast suggests that in the case of older infants (more than 12 months), this request was not simply linked to the need to be fed, but to a possible coping strategy for self-regulation in a stressful situation. These two coping strategies are efficient in providing self-comfort and limiting the negative effect of distress (51). This is in line with studies by Tronick (52), who stated that infant self-stimulation is a coping behavior that reduces stressful situations and offers self-calming, in particular when mothers are “still faced” (53). The mothers’ “absent look” is a kind of ecological “still face” situation. These interactive behaviors reflect the children’s coping strategies of self-regulation: they perceive these moments, and try to compensate with self-touch or by clinging to the satisfaction of primary needs. A previous research has shown that during a still-face procedure, when mothers were unavailable, infants spent more time touching themselves, supporting the regulatory and exploratory roles of infant touch, especially during periods of maternal unavailability (54).

It should be noted that, according to Main and Solomon “Indices of Disorganization and Disorientation (55), the children

did not show disorganized behavior during their mothers’ “absence,” but rather showed adaptive behaviors such as self-stimulation. Several explanations can be put forward for this. First, mothers had received psychological care from NGO psychosocial workers, which had possibly helped them to reduce their traumatic experience; our paradigm of trauma recall may perhaps not have reflected the fact that their trauma was reduced. Second, infant self-stimulation may be the first stage of coping; if the mother’s “absence” is repeated, their adaptive strategies may prove insufficient to handle the mother’s negative experience and may lead to internalized working models of disorganization. Third, the fact that the mothers talked very little to their child could limit the trauma transmission. Last, cultural differences might explain the way they cope with traumatic experiences. In traditional societies, children are not only raised by their mothers (56). They grow up surrounded by several caregivers, who complete the construction of the world and the child’s identity. This aspect of shared mothering could be a protective factor in the transmission of trauma from mother to child. The presence of a co-mothering system could mitigate or compensate for the effects of the mother’s lack of response to the child’s requests when she is absorbed by traumatic intrusions.

The third finding was that there were no significant associations between interaction and infant gender and age, the severity of traumatic experience assessed by IES, and mothers’ depression and anxiety symptoms assessed by HAD. This is maybe due to the small sample. Concerning age, it has however been shown (34) that the period when child locomotion develops might represent a critical time window for mother-child interaction in dyads with mothers having a history of abuse.

Not even the country of residence had an influence on the interactions between the mother and infant. The only difference was linked to the impact of the traumatic event and the dyads’ country of residence. The highest level of PTSD symptoms was associated with those living in the Central African Republic (CAR). This can be explained by the fact that people living in CAR are in an ongoing extremely traumatogenic context, whereas people who migrated and are living in refugee camps are more protected against new traumatizing situations.

Despite the fact that we found a significant positive association between the severity of the impact of the traumatic event (IES-R) and the time that had elapsed since the traumatic events, neither the severity of the traumatic experience nor the severity of anxiety or depression were associated with interactive behaviors. This may indicate that among the war traumatized and refugees, the theoretical frame to understand ITT has to be enlarged. Mothers’ general psychological distress, but not PTSD, was directly associated with negative parenting and child psychosocial difficulties in a sample of 291 Syrian refugee mothers in Lebanon who had been exposed to war trauma in the past (57). These results argue in favor of taking into account a broader framework for complex trauma in conflict and post-conflict areas, including the psychosocial framework, with the trauma focused approach. We did not evaluate the impact of other psychosocial conditions.

# LIMITATIONS OF THE STUDY

The main limitation of the study is the size of the sample. The recruitment in humanitarian contexts involves several challenges linked to security constraints and difficulties of access to subjects. A larger sample would be more representative of the observed mechanisms of transmission and make our results more significant. However, even with the small sample, our results point to particular lines of enquiry to pursue in future studies. The cultural particularities of mother and child relationships require investigation to design an appropriate framework for analysis as a key to understanding the affection linked to behaviors and interactions. The second main limitation is the absence of a control group. But our aim was to explore how trauma recall can affect mothers' presence to the child and the way they interact with the child, and not to globally qualify the interactive mother-child process. That is why we compared each dyad in three different settings, each dyad being its own control. A control group could have allowed us to compare mother-child interaction in play sessions. But by definition, it is not relevant to explore trauma recall with mothers who have not experienced traumatization. Another limitation is the context in which interaction took place, which was set up to catch the direct effects of trauma reactivation. The mothers were probably more involved in the interview than in playing with the child; however, this setting was close to an ecologic situation of daily life. Future studies should include longitudinal observation of mother and child dyads, with control groups, to identify the long-term impact of mothers' inappropriate response impacted by dissociation, when trauma is recalled, to their infants' needs and child development.

# CONCLUSION

This research is unique in its approach to mother-infant interactions: in the field explored (ITT in humanitarian contexts), by the setting (trauma recall), and by the microanalysis of interaction. Our results shed light on the process of trauma transmission, focusing on the caregiver reactions during trauma recall, and suggest that trauma affects a mother's availability to interact with her infant and to regulate his/her state of emotional arousal. The infant experiences this sudden lack of proper responses from the mother when she faces events or thoughts that trigger trauma memories, which can have an impact on the infant's perception of his/her own emotional status. A new and surprising finding was that children did not react by disorganization as has been previously described, but with self-stimulation, representing possible coping strategies in reaction to the mother's enigmatic reaction.

# REFERENCES

1. Beebe B, Lachmann F, Jaffe J, Markese S, Buck KA, Chen H, et al. Maternal postpartum depressive symptoms and 4-month mother-infant interaction. *Psychoanal Psychol.* (2012) 29:383–407. doi: 10.1037/a0029387

This adaptive behavior may compensate for the mothers' psychic absence and for the lack of "attunement" between mother and child.

Our results strongly reflect the increasing need to understand trauma transmission mechanisms in highrisk populations, particularly when they live in unremitting traumatogenic conditions.

In particular, the question about self-stimulation during mother's "absence" should be considered in further longitudinal studies as this initial adaptive mechanism could lead to internalized working model of disorganization, if repeated.

To prevent or to limit the impact of maternal trauma transmission, we recommend psychological interventions with mothers, starting from their pregnancy, and continuing after childbirth during the important period for the development of the child, until 2 years. Psychological support for at-risk mothers should aim to allow them to resolve the effects of the trauma in order to limit the transmission of trauma in interactions with the child. Interventions focusing on parental skills, their sensitivity, their ability to decipher, and respond to the specific needs of their child like video feed back interventions (58) are also recommended. But psychological support, which has been proposed for mothers with children born of sexual violence (59), has to go beyond trauma focused interventions, and provide a holistic and community embedded approach that can be applied across settings.

# DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.

# ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the institutional ethical review boards in CAR and in Cameroon, and by the Ministère de la Femme, de l'Action Sociale et de la Solidarité Nationale in Chad. Written informed consent to participate in this study was provided by the participants' legal guardian/next of kin.

# AUTHOR CONTRIBUTIONS

EDo was the principal investigator of this research and she was supervised by MF, MMo, CB, and LO during data collection and analysis. EDo, MMa, MMo, EDr, ML, and LO all contributed to the definition of the research methodology and contributed to the data analysis and interpretation. All authors participated in the writing and revision of the article.

2. Betancourt TS, McBain RK, Newnham EA, Brennan RT. The intergenerational impact of war: longitudinal relationships between caregiver and child mental health in postconflict sierra leone. *J Child Psychol Psychiatry.* (2015) 56:1101–7. doi: 10.1111/jcpp.12389
3. Roth M, Neuner F, Elbert T. Transgenerational consequences of PTSD: risk factors for the mental health of children whose mothers have



- been exposed to the Rwandan genocide. *Int J Ment Health Syst.* (2014) 8:12. doi: 10.1186/1752-4458-8-12
4. Field NP, Muong S, Sochanvimean V. Parental styles in the intergenerational transmission of trauma stemming from the khmer rouge regime in cambodia. *Am J Orthopsychiatry.* (2013) 83:483–94. doi: 10.1111/ajop.12057
5. Song SJ, Tol W, de Jong J. Indero: intergenerational trauma and resilience between burundian former child soldiers and their children. *Fam Process.* (2014) 53:239–51. doi: 10.1111/famp.12071
6. Bar-On D, Eland J, Kleber RJ, Krell R, Moore Y, Sagi A, et al. Multigenerational perspectives on coping with the holocaust experience: an attachment perspective for understanding the developmental sequelae of trauma across generations. *Int J Behav Dev.* (1998) 22:315–38. doi: 10.1080/016502598384397
7. Swenson C, Klingman A. Children and war. In: Saylor C, editor. *Children and Disasters.* New York, NY: Plenum Press (1993). p. 137–63. doi: 10.1007/978-1-4757-4766-9\_9
8. Ancharoff MR, Munroe JF, Fisher LM. The legacy of combat trauma: clinical implications of intergenerational transmission. In: Danieli Y, editor. *International Handbook of Multigenerational Legacies of Trauma.* New York, NY: Plenum Press (1998). p. 257–76. doi: 10.1007/978-1-4757-5567-1\_17
9. Coates S, Rosenthal J, Schechter D. (2003). *September 11: Trauma and Human Bonds. Relational Perspectives Book Series.* New York, NY: The Analytic Press/Taylor and Francis Group.
10. Schechter DS. Intergenerational communication of maternal violent trauma: understanding the interplay of reflective functioning and posttraumatic psychopathology. In: Coates SW, Rosenthal JL, Schechter DS, editors. *September 11: Trauma and Human Bonds.* Hillside, NJ: The Analytic Press Inc. (2003). p. 115–42.
11. Powers A, Stevens JS, O'Banion D, Stenson AF, Kaslow N, Jovanovic T, et al. *Intergenerational transmission of risk for PTSD symptoms in African American children: the roles of maternal and child emotion dysregulation.* *Psychol Trauma.* (2020). doi: 10.1037/tra0000543
12. Egeland B, Jacobvitz D, Sroufe LA. Breaking the cycle of abuse. *Child Dev.* (1988) 59:1080–1088. doi: 10.2307/1130274
13. Chamberlain C, Gee G, Harfield S, Campbell S, Brennan S, Clark Y, et al. Parenting after a history of childhood maltreatment: a scoping review and map of evidence in the perinatal period. *PLoS ONE.* (2019) 14:e0213460. doi: 10.1371/journal.pone.0213460
14. Kellermann NP. Transmission of holocaust trauma—an integrative view. *Psychiatry.* (2001) 64:256–67. doi: 10.1521/psyc.64.3.256.18464
15. Bowers ME, Yehuda R. Intergenerational transmission of stress in humans. *Neuropsychopharmacology.* (2016) 41:232–44. doi: 10.1038/npp.2015.247
16. Yehuda R, Lehrner A. Intergenerational transmission of trauma effects: putative role of epigenetic mechanisms. *World Psychiatry.* (2018) 17:243–57. doi: 10.1002/wps.20568
17. Lang AJ, Gartstein MA. Intergenerational transmission of traumatization: theoretical framework and implications for prevention. *J Trauma Dissoc.* (2018) 19:162–75. doi: 10.1080/15299732.2017.1329773
18. Bowlby J. *Attachment. Attachment and Loss Vol. I.* 2nd ed. New York, NY: Basic Book (1999).
19. Hesse E, Main M, Abrams KY, Rifkin A. Unresolved states regarding loss or abuse can have “second-generation” effects: disorganization, role-inversion, and frightening ideation in the offspring of traumatised non-maltreating parents. In: Solomon MF, Siegel DJ, editors. *Healing Trauma: Attachment, Mind, Body, Brain.* New York, NY: W. W. Norton and Co. (2003). p. 57–106.
20. Schuengel C, Bakermans-Kranenburg MJ, Van IJzendoorn MH. Frightening maternal behavior linking unresolved loss and disorganized infant attachment. *J Consult Clin Psychol.* (1999) 67:54–63. doi: 10.1037/0022-006X.67.1.54
21. Schechter DS, Coots T, Zeanah CH, Davies M, Coates SW, Trabka KA, et al. Maternal mental representations of the child in an inner-city clinical sample: Violence-related posttraumatic stress and reflective functioning. *Attachment Hum Dev.* (2005) 7:313–31. doi: 10.1080/14616730500246011
22. Lyons-Ruth K, Block D. The disturbed caregiving system: relations among childhood trauma, maternal caregiving, and infant affect and attachment. *Infant Ment Health J.* (1996) 17:77–95. doi: 10.1002/(SICI)1097-0355(199623)17:3<198::AID-IMHJ2>3.0.CO;2-L
23. Schechter DS, Coates SW, Kaminer T, Coots T, Zeanah CH, Davies M, et al. Distorted maternal mental representations and atypical behavior in a clinical sample of violence-exposed mothers and their toddlers. *J Trauma Dissoc.* (2008) 9:123–47. doi: 10.1080/15299730802045666
24. Aloni R, Mikulincer M, Zerach G, Solomon Z. The intergenerational sequelae of war captivity: the impact of a self-amplifying cycle of PTSD and attachment insecurities on offspring's attachment orientations. *Eur J Psychotraumatol.* (2020) 11:1741859. doi: 10.1080/2008198.2020.1741859
25. Biringen Z, Derscheid D, Vliegen N, Closson L, Easterbrooks MA. Emotional availability (EA): theoretical background, empirical research using the EA scales, and clinical applications. *Dev Rev.* (2014) 34:114–67. doi: 10.1016/j.dr.2014.01.002
26. van Ee E, Kleber RJ, Mooren TTM. War trauma lingers on: associations between maternal posttraumatic stress disorder, parent-child interaction, and child development. *Infant Ment Health J.* (2012) 33:459–68. doi: 10.1002/imhj.21324
27. Enlow MB, Egeland B, Carlson E, Blood E, Wright RJ. Mother-infant attachment and the intergenerational transmission of posttraumatic stress disorder. *Dev Psychopathol.* (2014) 26:41–65. doi: 10.1017/S0954579413000515
28. Schore AN. The effects of early relational trauma on right brain development, affect regulation, and infant mental health. *Infant Ment Health J.* (2001) 22:201–69. doi: 10.1002/1097-0355(200101/04)22:1<201::AID-IMHJ8>3.0.CO;2-9
29. Schore A. Dysregulation of the right brain: a fundamental mechanism of traumatic attachment and the psychopathogenesis of posttraumatic stress disorder (English). *Austr N Zeal J Psychiatry.* (2002) 36:9–30. doi: 10.1046/j.1440-1614.2002.00996.x
30. Stern D. *The Interpersonal World of the Infant: A View from Psychoanalysis and Developmental Psychology.* New York, NY: Basic Books (1985).
31. van Ee E, Kleber RJ, Jongmans MJ. Relational patterns between caregivers with PTSD and their nonexposed children: a review. *Trauma Violence Abuse.* (2016) 17:186–203. doi: 10.1177/1524838015584355
32. Lyons-Ruth K, Yellin C, Melnick S, Atwood G. Childhood experiences of trauma and loss have different relations to maternal unresolved and hostile-helpless states of mind on the AAI. *Attachment Hum Dev.* (2003) 5:330–52. doi: 10.1080/14616730310001633410
33. Sagi-Schwartz A, Van IJzendoorn MH, Grossmann KE, Joels T, Grossmann K, Scharf M, et al. Attachment and traumatic stress in female holocaust child survivors and their daughters. *Am J Psychiatry.* (2003) 160:1086–92. doi: 10.1176/appi.ajp.160.6.1086
34. Fuchs A, Möhler E, Resch F, Kaess M. Impact of a maternal history of childhood abuse on the development of mother-infant interaction during the first year of life. *Child Abuse Neglect.* (2015) 48:179–89. doi: 10.1016/j.chiabu.2015.05.023
35. Berthelot N, Ensink K, Bernazzani O, Normandin L, Luyten P, Fonagy P. Intergenerational transmission of attachment in abused and neglected mothers: the role of trauma-specific reflective functioning. *Infant Ment Health J.* (2015) 36:200–12. doi: 10.1002/imhj.21499
36. Dozio E, Feldman M, Moro M-R. Transmission du traumatisme mère-bébé dans les interactions précoces. *Pratiques Psychol.* (2016) 22:87–103. doi: 10.1016/j.prps.2016.01.001
37. Weiss DS, Marmar CR. The impact of event scale—revised. In: Wilson JP, Keane TM, Wilson JP, Keane TM, editors. *Assessing Psychological Trauma and PTSD.* New York, NY: Guilford Press (1997). p. 399–411. Retrieved from: <https://frodon.univ-paris5.fr/ur/!http://search.ebscohost.com/login.aspx?direct=true&db=psyhand&AN=1997-97162-013andlang=fr&site=ehost-live>
38. Joseph S. Psychometric evaluation of horowitz's impact of event scale: a review. *J Traumatic Stress.* (2000) 13:101–13. doi: 10.1023/A:1007777032063
39. Creamer M, Bell R, Failla S. Psychometric properties of the impact of event scale-revised. *Behav Res Ther.* (2003) 41:1489–96. doi: 10.1016/j.brat.2003.07.010
40. Weiss DS. The impact of event scale: revised. In: Wilson JP, Tang CS, Wilson JP, Tang CS, editors. *Cross-Cultural Assessment of Psychological Trauma and PTSD.* New York, NY: Springer Science + Business Media (2007). p. 219–238. Retrieved from: <https://frodon.univ-paris5.fr/ur/!http://search.ebscohost.com/login.aspx?direct=true&db=psyhand&AN=1997-97162-013andlang=fr&site=ehost-live>

- com/login.aspx?direct=true&db=psyhand&AN=2007-10108-010&lang=fr&site=ehost-live doi: 10.1007/978-0-387-70990-1\_10
41. Brunet A, St-Hilaire A, Jehel L, King S. Validation of a French version of the impact of event scale-revised. *Can J Psychiatry*. (2003) 48:56–61. doi: 10.1177/070674370304800111
  42. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand*. (1983) 67:361–70. doi: 10.1111/j.1600-0447.1983.tb09716.x
  43. Ali G-C, Ryan G, De Silva MJ. Validated screening tools for common mental disorders in low and middle income countries: a systematic review. *PLoS ONE*. (2016) 11:e0156939. doi: 10.1371/journal.pone.0156939
  44. Lebovici S, Mazet PH, Visier JP. *L'évaluation des Interactions Précoces Entre le Bébé et Ses Partenaires*. Paris: Eshel (1989).
  45. Beebe B, Jaffe J, Markese S, Buck K, Chen H, Cohen P, et al. The origins of 12-month attachment: a microanalysis of 4-month mother–infant interaction. *Attachment Hum Dev*. (2010) 12:3–141. doi: 10.1080/14616730903338985
  46. Sloetjes H, Wittenburg P. *Annotation by Category: ELAN and ISO DCR*. In: *LREC*. (2008) Retrieved from: [http://lrec-conf.org/proceedings/lrec2008/pdf/208\\_paper.pdf](http://lrec-conf.org/proceedings/lrec2008/pdf/208_paper.pdf)
  47. Scheff SW. Chapter 8 - nonparametric statistics. In: Scheff SW, editor. *Fundamental Statistical Principles for the Neurobiologist*. Academic Press (2016). p. 157–82. doi: 10.1016/B978-0-12-804753-8.00008-7
  48. Liotti G. Attachment and dissociation. *Dissoc Dissociat Disord*. (2009) 53–65.
  49. Pietromonaco PR, Barrett LF. The internal working models concept: what do we really know about the self in relation to others? *Rev Gen Psychol*. (2000) 4:155–75. doi: 10.1037/1089-2680.4.2.155
  50. Bailly L. Traumatisme psychique chez le jeune enfant et théories sociales infantiles. In: Baubet T, Lachal C, Ouss-Ryngaert L, Moro MR, editors. *Bébés et Traumas*. Grenoble: La Pensée sauvage (2006). p. 59–65.
  51. Koulomzin M, Beebe B, Anderson S, Jaffe J, Feldstein S, Crown C. Infant gaze, head, face and self-touch at 4 months differentiate secure vs. avoidant attachment at 1 year: a microanalytic approach. *Attachment Hum Dev*. (2002) 4:3–24. doi: 10.1080/14616730210123120
  52. Tronick EZ. Emotions and emotional communication in infants. *Am Psychol*. (1989) 44:112. doi: 10.1037/0003-066X.44.2.112
  53. Weinberg MK, Tronick EZ. Infant affective reactions to the resumption of maternal interaction after the still-face. *Child Dev*. (1996) 67:905–14. doi: 10.2307/1131869
  54. Moszkowski RJ, Stack DM. Infant touching behaviour during mother–infant face-to-face interactions. *Infant Child Dev*. (2007) 16:307–19. doi: 10.1002/icd.510
  55. Main M, Solomon J. *Indices of Disorganization and Disorientation* (1990).
  56. Moro MR. *Psychothérapie transculturelle des enfants de migrants*. Dunod (1998).
  57. Sim A, Bowes L, Gardner F. Modeling the effects of war exposure and daily stressors on maternal mental health, parenting, and child psychosocial adjustment: a cross-sectional study with Syrian refugees in Lebanon. *Glob Ment Health*. (2018) 5:e40. doi: 10.1017/gmh.2018.33
  58. Schechter DS, Moser DA, Reliford A, McCaw JE, Coates SW, Turner JB, et al. Negative and distorted attributions towards child, self, and primary attachment figure among posttraumatically stressed mothers: what changes with clinician assisted videofeedback exposure sessions (CAVES). *Child Psychiatry Hum Dev*. (2015) 46:10–20. doi: 10.1007/s10578-014-0447-5
  59. Anderson DI, Campos JJ, Witherington DC, Dahl A, Rivera M, He M, et al. The role of locomotion in psychological development. *Front Psychol*. (2013) 4:440. doi: 10.3389/fpsyg.2013.00440
  60. Murray L, Fiori-Cowley A, Hooper R, Cooper P. The impact of postnatal depression and associated adversity on early mother–infant interactions and later infant outcome. *Child Dev*. (1996) 67:2512–26. doi: 10.2307/1131637
  61. Lovejoy MC, Graczyk PA, O'Hare E, Neuman G. Maternal depression and parenting behavior. *Clin Psychol Rev*. (2000) 20:561–92. doi: 10.1016/S0272-7358(98)00100-7
  62. Stern D. *The Present Moment in Psychotherapy and Everyday Life*. New York, NY: W.W. Norton and Company (Norton Series on Interpersonal Neurobiology) (2004).
  63. Schreier A, Wittchen H-U, Hofler M, Lieb R. Anxiety disorders in mothers and their children: prospective longitudinal community study. *Br J Psychiatry*. (2008) 192:308–9. doi: 10.1192/bjp.bp.106.033589

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Dozio, Feldman, Bizouerne, Drain, Laroche Joubert, Mansouri, Moro and Ouss. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.